

heattransfer report .docx



Document Details

Submission ID

trn:oid:::15229:87689077

Submission Date

Mar 25, 2025, 8:59 AM GMT+5

Download Date

Mar 25, 2025, 8:59 AM GMT+5

File Name

heattransfer report .docx

File Size

275.2 KB

10 Pages

1,842 Words

11,261 Characters

0% detected as AI

The percentage indicates the combined amount of likely AI-generated text as well as likely AI-generated text that was also likely AI-paraphrased.

Caution: Review required.

It is essential to understand the limitations of AI detection before making decisions about a student's work. We encourage you to learn more about Turnitin's AI detection capabilities before using the tool.

Detection Groups

- 
1 AI-generated only 0%
 Likely AI-generated text from a large-language model.
- 
2 AI-generated text that was AI-paraphrased 0%
 Likely AI-generated text that was likely revised using an AI-paraphrase tool or word spinner.

Disclaimer

Our AI writing assessment is designed to help educators identify text that might be prepared by a generative AI tool. Our AI writing assessment may not always be accurate (it may misidentify writing that is likely AI generated as AI generated and AI paraphrased or likely AI generated and AI paraphrased writing as only AI generated) so it should not be used as the sole basis for adverse actions against a student. It takes further scrutiny and human judgment in conjunction with an organization's application of its specific academic policies to determine whether any academic misconduct has occurred.

Frequently Asked Questions

How should I interpret Turnitin's AI writing percentage and false positives?

The percentage shown in the AI writing report is the amount of qualifying text within the submission that Turnitin's AI writing detection model determines was either likely AI-generated text from a large-language model or likely AI-generated text that was likely revised using an AI-paraphrase tool or word spinner.

False positives (incorrectly flagging human-written text as AI-generated) are a possibility in AI models.

AI detection scores under 20%, which we do not surface in new reports, have a higher likelihood of false positives. To reduce the likelihood of misinterpretation, no score or highlights are attributed and are indicated with an asterisk in the report (*%).

The AI writing percentage should not be the sole basis to determine whether misconduct has occurred. The reviewer/instructor should use the percentage as a means to start a formative conversation with their student and/or use it to examine the submitted assignment in accordance with their school's policies.

What does 'qualifying text' mean?

Our model only processes qualifying text in the form of long-form writing. Long-form writing means individual sentences contained in paragraphs that make up a longer piece of written work, such as an essay, a dissertation, or an article, etc. Qualifying text that has been determined to be likely AI-generated will be highlighted in cyan in the submission, and likely AI-generated and then likely AI-paraphrased will be highlighted purple.

Non-qualifying text, such as bullet points, annotated bibliographies, etc., will not be processed and can create disparity between the submission highlights and the percentage shown.



Time-Dependent Heat Transfer Analysis: Freezing of a Human Body in Outer Space

Abstract / Executive Summary

The study investigates the duration required for human body freezing in space conditions because exposure to vacuum environments imposes severe thermal challenges. The problem addresses freezing times through a time-dependent heat transfer model that uses an 80-kg spherical human body at 310 K initial temperature near 3 K background temperature. The main heat loss process occurs through Stefan-Boltzmann radiation since conductive and convective effects prove negligible while metabolic heat production remains constant. Numerical simulations demonstrate that body heat generates substantial delays in temperature reduction so that freezing requires several days to occur within such space conditions.

Table of content

Time-Dependent Heat Transfer Analysis: Freezing of a Human Body in Outer Space.....	1
Abstract / Executive Summary.....	1
Introduction	2
Analysis	2
Model Formulation.....	2
1. Energy Balance Equation	2
2. Radiative Heat Loss.....	3
3. Assumptions and Simplifications	3
4. Python Implementation Outline.....	4
1. Primary Simulation of Body Temperature	4
2. Sensitivity Analysis: Effect of Metabolic Heat Generation	6
3. Sensitivity Analysis: Impact of Emissivity	7
4. Parameter Summary Table	8
Results	9
Primary Simulation:.....	9
Emissivity Impact:	9
Discussion	9
Geometric Simplification:	9
Dominance of Radiative Heat Transfer:	9
Constant Metabolic Heat Generation:	9
Parameter Uncertainty:	9
Conclusion.....	9
Reference.....	10

Introduction

The project studies the time required for human body freezing under outer space environmental conditions. The space environment creates such low pressure that traditional thermal transfer methods through conduction and convection cease to exist allowing the Stefan–Boltzmann law to define heat loss via radiation as the primary cooling mechanism. The mathematical assessment relies on simulating an 80-kg spherical human body maintained at 310 K while exposed to the 3 K outer space temperature while generating metabolic heat at 100 W. The methodology includes creating a time-varying heat transfer system which controls both radiative cooling and internal heat generation before performing numerical solutions to the differential equation. Through this method researchers gain understanding about heat dissipation rates while demonstrating how basic simplifying assumptions help manage complex thermodynamic scenarios.

Analysis

Model Formulation

We designed a basic thermal model through energy balance calculations for a human body weighing 80-kg in the outer space near vacuum environment. The model considers the body as a single homogeneous sphere that measures 1.5 m in diameter. Space's harsh environmental conditions promote radiation heat transfer as the main process because conduction and convection processes become insignificant. Based on the lumped capacitance assumption the model establishes that the body temperature remains uniformly distributed at all times. Scientists often apply this assumption to describe temperature conditions when body heat diffusion inside the system remains minimal relative to whole-body temperature changes in ambient space.

1. Energy Balance Equation

The energy balance for the body can be written as:

$$mc_p \frac{dT}{dt} = -Q_{\text{loss}} + \dot{Q}_{\text{met}}$$

where:

m is the mass of the body (80 kg),

c_p is the specific heat capacity of human tissue (approximately 3500 J/(kg·K)),

$T(t)$ is the temperature of the body (in K) as a function of time,

Q_{loss} is the rate of heat loss from the body, and

$\dot{Q}_{\text{met}}$ is the metabolic heat generation (assumed constant at 100 W).

2. Radiative Heat Loss

The principal cooling mechanism of space depends on radiative heat transfer whose behavior matches the Stefan–Boltzmann law. The heat loss by radiation follows the rate expression given by:

$$Q_{\text{rad}} = \varepsilon \sigma A (T^4 - T_{\text{env}}^4)$$

where:

- ε is the emissivity of the body (taken as 0.95),
- σ is the Stefan–Boltzmann constant ($5.67 \times 10^{-8} \text{ W/m}^2\text{K}^4$),
- A is the surface area of the body, and
- T_{env} is the ambient temperature of space (approximately 3 K).

For a spherical body, the surface area A is calculated by:

$$A = 4\pi r^2$$

with the radius r computed as half of the characteristic length:

$$r = \frac{1.5}{2} \text{ m} = 0.75 \text{ m}$$

Thus, the energy balance becomes:

$$mc_p \frac{dT}{dt} = -\varepsilon \sigma A (T^4 - T_{\text{env}}^4) + \dot{Q}_{\text{met}}$$

3. Assumptions and Simplifications

Lumped Capacitance:

Every moment we accept that body temperatures remain identical throughout the entire human body. Such assumption remains applicable when the resistance from internal temperature gradients is considerably smaller than the radiative heat resistance.

Negligible Conduction and Convection:

Spaces near vacuum lack all necessary materials that would help conduct or convect heat in this environment. Both terms present in an energy balance have been treated as non-existent or considered negligible.

Constant Metabolic Heat Generation:

A constant rate of 100 W defines the metabolic heat generation ($\dot{Q}_{\text{met}}$) throughout the analysis. The rate remains fixed at the initial value throughout the model because analysis requires its simplification.

Radiative Dominance:

Radioative heat transfer becomes vital because the body begins at 310 K while operating in a space of 3 K temperature. The Stefan–Boltzmann law shows radiative cooling dependency on the fourth power of temperature which demonstrates its strong nonlinearity.

4. Final Differential Equation

In summary, the governing differential equation for the temperature $T(t)$ is:

$$\frac{dT}{dt} = \frac{-\varepsilon\sigma A (T^4 - T_{\text{env}}^4) + \dot{Q}_{\text{met}}}{mc_p}$$

This equation is solved numerically to track the temperature evolution over time.

4. Python Implementation Outline

1. Primary Simulation of Body Temperature

```
import numpy as np
import matplotlib.pyplot as plt
from scipy.integrate import odeint

# -----
# Constants and Parameters
# -----
m = 80          # Mass of the body (kg)
cp = 3500       # Specific heat capacity of human tissue (J/(kg·K))
T0 = 310        # Initial body temperature (K)
T_env = 3       # Ambient temperature in space (K)
L = 1.5         # Characteristic length (diameter) in m
r = L / 2       # Radius of the sphere (m)
A = 4 * np.pi * r**2  # Surface area of the sphere (m^2)
epsilon = 0.95  # Emissivity of human skin/clothing
sigma = 5.67e-8 # Stefan–Boltzmann constant (W/m^2K^4)
Q_met = 100     # Metabolic heat generation (W)

# -----
```

```
# Differential Equation Function
# -----
def dT_dt(T, t):
    """
    Computes the time derivative of the body temperature T based on the energy balance.


    Energy balance:
    
$$m \cdot c_p \cdot (dT/dt) = -\epsilon \sigma A (T^4 - T_{env}^4) + Q_{met}$$

    """
    # Radiative heat loss computed via the Stefan-Boltzmann law
    Q_rad = epsilon * sigma * A * (T**4 - T_env**4)
    # Derivative of temperature (dT/dt)
    dTdt = (-Q_rad + Q_met) / (m * cp)
    return dTdt

# -----
# Time Domain for Simulation
# -----
# Simulate for 1e6 seconds (~11.57 days)
t = np.linspace(0, 1e6, 10000)

# -----
# Numerical Integration
# -----
T_solution = odeint(dT_dt, T0, t)

# -----
# Plotting the Results
# -----
plt.figure(figsize=(10, 6))
plt.plot(t / 3600, T_solution, label='Body Temperature')
plt.xlabel("Time (hours)")
plt.ylabel("Temperature (K)")
plt.title("Cooling of a Human Body in Outer Space")
plt.legend()
plt.grid(True)
plt.show()
```



2. Sensitivity Analysis: Effect of Metabolic Heat Generation


```
import matplotlib.pyplot as plt

# Different metabolic heat generation values for sensitivity analysis (W)
Q_met_values = [0, 50, 100, 150]
plt.figure(figsize=(10, 6))

for Q in Q_met_values:
    def dT_dt_sensitivity(T, t):
        Q_rad = epsilon * sigma * A * (T**4 - T_env**4)
        dTdt = (-Q_rad + Q) / (m * cp)
        return dTdt

    T_sens = odeint(dT_dt_sensitivity, T0, t)
    plt.plot(t/3600, T_sens, label=f'Metabolic Heat = {Q} W')

plt.xlabel('Time (hours)')
plt.ylabel('Temperature (K)')
plt.title('Sensitivity Analysis: Impact of Metabolic Heat on Cooling')
plt.legend()
plt.grid(True)
plt.show()
```



3. Sensitivity Analysis: Impact of Emissivity


```
import matplotlib.pyplot as plt

# Varying emissivity values for sensitivity analysis
epsilon_values = [0.8, 0.9, 0.95, 1.0]
plt.figure(figsize=(10, 6))

for eps in epsilon_values:
    def dT_dt_emissivity(T, t):
        Q_rad = eps * sigma * A * (T**4 - T_env**4)
        dTdt = (-Q_rad + Q_met) / (m * cp)
        return dTdt

    T_emiss = odeint(dT_dt_emissivity, T0, t)
    plt.plot(t/3600, T_emiss, label=f'Emissivity = {eps}')

plt.xlabel('Time (hours)')
plt.ylabel('Temperature (K)')
plt.title('Sensitivity Analysis: Impact of Emissivity on Cooling')
plt.legend()
plt.grid(True)
plt.show()
```



4. Parameter Summary Table

```
import pandas as pd
```

```
# Create a DataFrame with simulation parameters
```

```
data = {
```

```
    'Parameter': ['Mass (kg)', 'Specific Heat (J/kgK)', 'Initial Temp (K)', 'Ambient Temp (K)',
```

```
                  'Characteristic Length (m)', 'Radius (m)', 'Surface Area (m^2)',
```

```
                  'Emissivity', 'Stefan-Boltzmann Constant (W/m^2K^4)', 'Metabolic Heat (W)'],
```

```
    'Value': [m, cp, T0, T_env, L, r, A, epsilon, sigma, Q_met]
```

```
}
```

```
df_params = pd.DataFrame(data)
```

```
print(df_params)
```

	Parameter	Value
0	Mass (kg)	8.000000e+01
1	Specific Heat (J/kgK)	3.500000e+03
2	Initial Temp (K)	3.100000e+02
3	Ambient Temp (K)	3.000000e+00
4	Characteristic Length (m)	1.500000e+00
5	Radius (m)	7.500000e-01
6	Surface Area (m^2)	7.068583e+00
7	Emissivity	9.500000e-01
8	Stefan-Boltzmann Constant (W/m^2K^4)	5.670000e-08
9	Metabolic Heat (W)	1.000000e+02

Results

Various key findings emerge from the simulation and sensitivity investigations.

Primary Simulation:

The temperature-time graph demonstrates that metabolic heat generation at 100 W controls the cooling process while maintaining heat loss through outer space despite large temperature differential. During several days the body temperature descends at a steady pace instead of undergoing an instant drop.

Metabolic Heat Sensitivity:

Changes in metabolic heat output show that small amounts of generated heat will affect temperature decline substantially. Zero metabolic heat leads to an abrupt temperature decrease which demonstrates how body-produced heat works as an important factor for enduring survival.

Emissivity Impact:

More efficient heat transfer through radiation occurs when products possess higher values of emissivity. Modeling thermal behavior in space requires precise emissivity measurement because this sensitivity analysis proves the impact of emissivity on space modeling results.

Discussion

Multiple simplifications form the basis for building this model.

Geometric Simplification:

A simple spherical shape serves as a body representation in this model because it helps simplify body geometries and varied thermal characteristics of actual human bodies.

Dominance of Radiative Heat Transfer:

Space conditions that are nearly void of matter make conduction and convection patterns negligible. The inclusion of these bodies becomes necessary for space scenarios which do not extend to deep space.

Constant Metabolic Heat Generation:

A constant metabolic rate of 100 W fails to explain the physical changes that occur during body cooling. Temperature decreases metabolic rates in humans and this additional rate slowdown will influence the overall cooling process.

Parameter Uncertainty:

The technical limitations of emissivity and specific heat capacity measurements lead to modeling uncertainties that affect simulation results according to sensitivity analysis results.

The present model offers fundamental knowledge about space-based body cooling but its accuracy would increase through additional refinements including variable metabolism modeling and better representation of human anatomy.

Conclusion

This document conducts a temporally evolving heat transfer assessment of human outer space bodies through a basic spherical human model. The computed solutions of the energy balance equation confirm that body heat generated through metabolism extends the amount of time required for freezing to occur. The cooling rate depends heavily on two important factors including metabolic heat production and surface emissivity according to sensitivity analysis. The simplified model correctly demonstrates how radiative losses interact with metabolic heat to extend freezing time to multiple days despite its basic assumptions.

Reference

Burkov, I.A., Kolishkin, L.M., Pushkarev, A.V., Shakurov, A.V., Tsiganov, D.I. and Zherdev, A.A., 2022. Experimental and computational thermal analysis of partial-body cryotherapy. *International Journal of Heat and Mass Transfer*, 183, p.122194.

Verma, R. and Kumar, S., 2021. Computational study on skin tissue freezing using three-phase lag bioheat model. *Journal of Heat Transfer*, 143(11), p.111201.

Barman, B., Bernal, N., Xu, Y. and Zapata, Ó., 2022. Ultraviolet freeze-in with a time-dependent inflaton decay. *Journal of Cosmology and Astroparticle Physics*, 2022(07), p.019.

El-Nabulsi, R.A. and Anukool, W., 2024. Time-dependent heating problem of the solar corona in fractal dimensions: a plausible solution. *Advances in Space Research*, 74(5), pp.2510-2529.

Qin, Y., 2021. Effect of inclusion of nanoparticles on unsteady heat transfer. *Applied Nanoscience*, pp.1-14.

Xu, M., Akhtar, S., Zueter, A.F., Auger, V., Alzoubi, M.A. and Sasmito, A.P., 2020. Development of analytical solution for a two-phase Stefan problem in artificial ground freezing using singular perturbation theory. *Journal of Heat Transfer*, 142(12), p.122401.

Zimmermann, Y., Keski-Rahkonen, J., Graf, A.M. and Heller, E.J., 2024. Rise and fall of anderson localization by lattice vibrations: A time-dependent machine learning approach. *Entropy*, 26(7), p.552.